arXiv:1202.1230v1 [quant-ph] 6 Feb 2012

On-chip past-future quantum correlations

Carlos Sabín,[1] Borja Peropadre,[1] Marco del Rey,[1] and Eduardo Martín-Martínez[2]
[1] *Instituto de Física Fundamental, CSIC, Serrano 113-B, 28006 Madrid, Spain*[*]
[2] *Institute for Quantum Computing, University of Waterloo,*
200 Univ. Avenue W, Waterloo, Ontario N2L 3G1, Canada

(Dated: February 7, 2012)

We propose a realistic circuit QED experiment to test the extraction of past-future vacuum entanglement to a pair of superconducting qubits. The qubit P interacts with the quantum field along an open transmission line for an interval $T_{\rm on}$ and then, after a time-lapse $T_{\rm off}$, the qubit F starts interacting for a time $T_{\rm on}$ in a symmetric fashion. After that, past-future quantum correlations of the vacuum will have transferred to the qubits, even if the qubits do not coexist at the same time. We show that this experiment can be realized with current technology and discuss its utility as a possible implementation of a quantum memory.

PACS numbers: 42.50.-p, 03.65.Ud, 03.67.Lx, 85.25.-j

The fact that the vacuum of a quantum field is an entangled state was discovered long ago [1–3] , but it was considered a mere formal result until it was addressed from an applied perspective in [4]. Since then, this intriguing property has attracted a great deal of attention as a possible new resource for Quantum Information tasks [5–8].

As shown in [4], the entanglement contained in the vacuum of a scalar field can be transferred to a pair of two-level spacelike separated detectors interacting with the field at the same time. Unfortunately, this theoretical result seems to be very difficult from an experimental viewpoint, even in the context of a trapped-ion simulation [5]. Recently, it has also been shown [9] that the vacuum of a massless scalar field contains quantum correlations [10] between the future and the past light cones. A theoretical method of extraction by transference to detectors interacting with the field at different times has also been proposed [11] using detectors with time-dependent energy gaps whose experimental implementation seems extremely challenging. Another ideal proposal involving cavities transparent to a single mode was provided in [12], although this setting seems even more difficult to tackle experimentally.

On the other hand, circuit QED [13] provides a framework in which the interaction of two-level systems with a quantum field can be naturally considered. The combination of superconducting qubits with transmission lines implement an artificial 1-D matter-radiation interaction, with the advantage of a large experimental accessibility and tunability of the physical parameters. Using these features, fundamental problems in Quantum Field Theory hitherto considered as ideal are now accessible to experiment [14]. In particular, the possibility of achieving an ultrastrong coupling regime [15–17] has already been exploited to propose a feasible experimental test of the extraction of vacuum entanglement to a pair of spacelike separated qubits [7].



FIG. 1: Experimental proposal for past-future entanglement extraction. a) Time evolution of our protocol: the qubit P, initially prepared in its excited state, interacts with the vacuum field ($\Delta\psi$) for a time T_{on}. After a certain time $T_{\rm off}$ with no interaction, a second qubit F interacts with the field getting entangled with the qubit P. b) Scheme leading to a switchable coupling between a qubit and the microwave quantum field.

In this work we will take advantage of the aforementioned features of circuit QED in the ultrastrong coupling regime in order to propose a realistic experiment for the extraction of past-future correlations [18] contained in the vacuum of a quantum field. We will consider a setup consisting of a pair of superconducting qubits P and F separated by a fixed distance r in a common open transmission line (Fig. 1a). First, the interaction of P with the vacuum of the field, which is tunable in strength, is on for a time interval $T_{\rm on}$ - the past. Then, P is disconnected from the field during a given $T_{\rm off}$. Finally, the interaction of F is switched on during $T_{\rm on}$ -the future - while keeping P disconnected. After this procedure, we will show that the qubits end up in a strongly correlated quantum state, in spite of the fact that they have never interacted with the field at the same time. Remarkably, the correlations between different times contained in the vacuum of the quantum field have been transferred to the qubits. We will thoroughly discuss the relationship of this effect with the spacetime region at which the qubits are placed, or in other words, with r, the times $T_{\rm on}$, $T_{\rm off}$ and the propagation velocity of the field v. We stress that our proposal is free of idealized requirements such as complicated time-dependent gaps and fully within reach of current circuit

QED technologies.

In addition to its interest from the fundamental viewpoint, our protocol has also an important applied counterpart. As suggested in [11], the extraction of past-future quantum correlations enables its use as a quantum channel for quantum teleportation "in time". We will show how this opens the door to a novel kind of quantum memory in which the information of the quantum state of some ancillary qubit P' is codified in the field during $T_{\rm off}$ and then recovered in F using classical information stored in the past - even if P is destroyed after its interaction with the field.

Let us explain the formalism and our results in more detail. From now on we focus on a setup of circuit-QED with two superconducting qubits P and F interacting via a quantum field. The qubits have two stationary states $|e\rangle$ and $|g\rangle$ separated by an energy $\hbar\Omega$ and might interact with a one-dimensional field, $V(x)$, which propagates along a one-dimensional open microwave guide or transmission line that connects them

$$V(x) = i\int_{-\infty}^{\infty} dk\,\sqrt{N\omega_k}\,e^{ikx}a_k + \text{H.c.} \tag{1}$$

This field has a continuum of Fock operators $[a_k, a^\dagger_{k'}] = \delta(k-k')$, and a linear spectrum, $\omega_k = v|k|$, where v is the propagation velocity of the field. The normalization N and the speed of photons, $v = (cl)^{-1/2}$, depend on the microscopic details such as the capacitance and inductance per unit length, c and l. We will assume qubits that are much smaller than the relevant wavelengths, $\lambda = 2\pi\, v/\Omega_J$, $(J = P, F)$ and are separated by a fixed distance r. Under these conditions the Hamiltonian, $H = H_0 + H_I$, splits into a free part for the qubits and the field $H_0 = \frac{1}{2}\hbar(\Omega_P\sigma^z_P + \Omega_F\sigma^z_F) + \int_{-\infty}^{\infty} dk\,\hbar\omega_k a^\dagger_k a_k$ and a point-like interaction between them

$$H_I = -\sum_{J=P,F} d_J\, V(x_J)\,\sigma^x_J = H_{IP} + H_{IF} \tag{2}$$

Here x_J are the fixed positions of the atoms, and $d_J\,\sigma^x_J$ is a dimensional reduction of the matter- radiation interaction hamiltonian with two-level atoms and the electromagnetic field, analogous - but not fully equivalent - to the Unruh-de Witt model. [19].

Representing the states in terms of qubit (P, F) and field (V) free eigenstates with the notation $|\psi\rangle = |PFV\rangle$, the initial state would be $|\psi(-t_2)\rangle = |eg0\rangle$, where only qubit P has been excited, while both F and the field V remain in their ground and vacuum states, respectively. According to our past-future scheme (Fig. 1a), the system evolves in the interaction picture into the state

$$|\psi(t_2)\rangle = \mathcal{T} e^{-i\int_{-t_2}^{t_2}\frac{dt'}{\hbar}[\Theta(-t'-t_1)H_{IP}^{(t')}+\Theta(t'-t_1)H_{IF}^{(t')}]}\,|eg0\rangle\,, \tag{3}$$

$\mathcal{T}$ being the time ordering operator.

We use the formalism of perturbation theory up to the second order and beyond Rotating Wave Approximation [7] and trace over the field degrees of freedom to obtain the corresponding two-qubit reduced density matrix ρ_{PF} evaluated at t_2. The degree of entanglement of this X-state can be characterized with the concurrence, which at the considered order is given by:

$$\mathcal{C}(\rho_{PF}) = 2\left(|X| - \left(\sum_k |A_{1,k}|^2\right)^{1/2}\left(\sum_k |B_{1,k}|^2\right)^{1/2}\right), \tag{4}$$

X standing for the normalized amplitude of photon - real and virtual - exchange and $\sum_k |A_{1,k}|^2$, $\sum_k |B_{1,k}|^2$ for the total normalized probability of single-photon emission by qubit P and F, respectively. These terms can be computed analytically - following similar techniques as in [7] - as a function of four dimensionless parameters, $\xi_{\rm on}$, $\xi_{\rm off}$, K_P and K_F. The first two, $\xi_{\rm on} = vT_{\rm on}/r$, $\xi_{\rm off} = vT_{\rm off}/r$ allow to discriminate the different spacetime regions. The remaining ones are dimensionless coupling strengths for qubits P and F: $K_J = \frac{4d_J^2 N}{\hbar^2 v} = 2\left(\frac{g_J}{\Omega_J}\right)^2$. We will restrict to consider times where $2K_J\Omega_J t_2 \ll 1$, needed for our perturbative approach to remain valid.

Three different regions emerge from the parameters above: if $T_{\rm on} + T_{\rm off} > r/v$ (we will refer to this spacetime region as region 3), F may start at $t = t_1$ to absorb radiation emitted by P in the past (if $T_{\rm off} < r/v$) or not absorb radiation at all (if $T_{\rm off} > r/v$). If $2T_{\rm on} + T_{\rm off} > r/v$ but $T_{\rm on} + T_{\rm off} < r/v$ (region 2), F will start to absorb radiation somewhen in the future. If $2T_{\rm on} + T_{\rm off} < r/v$ (region 1), there is no real photon exchange, but vacuum correlations - or virtual photon exchange - are allowed at any time. Only in region 1 and in region 3 for $T_{\rm off} > r/v$ can it be said that we are dealing with a pure effect of transference of the quantum correlations between the past and the future contained in the vacuum. In the other two spacetime regions these correlations are assisted by the photon exchange, operating at any time in the future - region 3 for $T_{\rm off} < r/v$ - or only for a given time - region 2 -.

In Fig. 2a we sketch the behaviour of the concurrence as a function of $T_{\rm on}$ and $T_{\rm off}$, for coupling strengths $g_J/\Omega_J \simeq 0.1$, such as in cutting-edge experiments of ultrastrong coupling in circuit QED [16, 17] and accessible values of the qubit's gaps and distance. We note that qubit-qubit entanglement is sizable in the ns regime, both in the previously defined regions 2 and 3 of the spacetime. In addition, Fig. 2b shows a significant amount of entanglement in region 1, entailing a pure transference of vacuum correlations. In general, entanglement is concentrated around $\Omega_J\, T_{\rm on} \simeq \Omega_J\, T_{\rm off} \simeq 1$ and $\xi_{\rm on} \simeq \xi_{\rm off} \simeq 1$. Thus, for qubit distances of the order of λ, entanglement shows up in the ns regime, but drifts towards shorter times as the distance diminishes, as can be seen in Fig. 2c.

In particular, in our analysis with time independent energy gaps, concurrence is 0 in the purely time-like scenario where $T_{\rm off} > r/v$, including the $r = 0$ case, as



FIG. 2: a) Concurrence vs. $T_{\rm on}$ and $T_{\rm off}$ for $g = g_P = g_F$, $\Omega = \Omega_P = \Omega_F = 2\pi \times 1\,\mathrm{GHz}$, $g/\Omega = \sqrt{3}/2 \cdot 10^{-1}$ and $r/\lambda = 2$. Significant entanglement is generated at both sides of the line $T_{\rm on} + T_{\rm off} = 2\,\mathrm{ns}$ which discriminates regions 2 and 3. b) Concurrence vs $T_{\rm on}$ and the coupling strength g/Ω for $T_{\rm off} = 0$, $\Omega = \Omega_P = \Omega_F = 2\pi \times 1\,GHz$, $r/\lambda = 2$, $g = g_P = g_F$. Sizable entanglement is generated at both sides of the line $T_{\rm on} = 1\,\mathrm{ns}$ which discriminate regions 1 and 2. c) Concurrence vs. $T_{\rm off}$ and r/λ for $T_{\rm on} = 1\,\mathrm{ns}$, $g = g_P = g_F$, $\Omega = \Omega_P = \Omega_F = 2\pi \times 1\,\mathrm{GHz}$, and $g/\Omega = \sqrt{3}/2 \cdot 10^{-1}$. The generation of entanglement is concentrated around $(r/\lambda)/1\,\mathrm{GHz} \simeq (T_{\rm on} + T_{\rm off})$.

described in [11]. As mentioned before, a tailor-made time-dependence for the qubit gap ($\propto 1/t$) would be required in order to access vacuum quantum correlations in such a scheme. As a matter of fact, the proposal in [11] exploits a formal analogy [9]- only fulfilled for massless fields - between the past and the future light cones and the left-right Rindler wedges. However, one must be very careful about the extent to what this analogy is valid: while it is possible to think of the vacuum state as an entangled state of modes observed by causally disconnected observers in the left-right wedges of the space-time [20], it is not clear whether this way of thinking can be transported to the past-future light cones case [21]. Thus, standard detectors do not actually couple to the entangled modes of the past and the future light cones and the time dependent gap must be introduced. On the experimental side, one could think of engineering a tunable gap with charge qubits, but in this case the qubit-field interaction could not be connected and disconnected at will. While the latter condition can be indeed circumvented with our setup, the $1/t$ dependence would result extremely challenging from the experimental viewpoint.

Given this, we will focus on a more feasible setting, aiming to test the results shown in Fig. 2. As mentioned before, it consists of a circuit QED design where two superconducting qubits, galvanically coupled to the center conductor of an open transmission line, interact with the vacuum field, being this interaction tunable in strength [22] (see Fig. 1a). A first superconducting qubit P (prepared in its excited state by driving it with a microwave pulse) interacts for a time $T_{\rm on}$ with the vacuum field. Subsequently, we switch off the interaction by applying an external flux bias (Fig. 1b). Finally, after a time $T_{\rm off}$, we switch on the interaction of a second qubit F, generating a certain amount of entanglement between P and F, which can be quantified with quantum state tomography [23]. It must be noted that, even if what we calculate here is a standard entanglement measure (namely the concurrence for qubits P and F at time $t = t_2$), we could still use this quantity as a measure of quantum correlations between past and future for the case where qubit P had been destroyed after $t = -t_1$. In this case, the only novelty of the tomography process will be that the local measurements with correlated outcomes would occur at different times.

So far we have introduced a qualitative scheme based on switchable interactions. This key ingredient can be engineered using the design depicted in Fig. 1b. Intuitively the system can be seen as a flux qubit (top ring) coupled to the transmission line (vacuum field) by means of the intermediate loop, which regulates the coupling by varying the external flux bias (Φ_3) passing through the SQUID. Actually the intermediate loop is part of the qubit and part of the field, and as such it has to be quantized together with them. Only then qubit, tunable coupling and quantum field would emerge as effective degrees of freedom.

Since we remain in the flux regime ($E_C \ll E_J$) we can just consider the dominant contributions to the energy given by the Josephson junctions, leading to the effective Hamiltonian $H_{\rm eff} = -\sum_{i=1}^{5} E_{J_i} \cos(\varphi_i)$, where E_{J_i} and φ_i stand for the Josephson energy and gauge - invariant phase of the i-th junction respectively. The flux quantization over closed loops imposes the following restrictions, $\sum_k \varphi_k = f_k \pmod{2\pi}$, where the parameters $f_k = 2\pi\Phi_k/\Phi_0$ allow us to select the qubit gap

Ω_J, the shape of the interaction, and most important, the strength of the latter. Considering the standard flux qubit configurations ($E_{J1} = E_{J1} = \alpha E_{J3}$), together with $E_{J4} = E_{J5} = \alpha_4 E_{J1}$, the effective Hamiltonian can be decomposed as the free Hamiltonians of the qubit and the electromagnetic field with a certain interaction given by $H_{\text{int}}/E_{J1} = -\int dk \alpha_4 \cos(f_3)\, \sigma_x \Delta\psi$, where f_3 stands for the normalized external flux bias passing through the SQUID, and $\Delta\psi \propto (a_k + a_k^\dagger)$. It is now clear how by driving an external flux $\Phi_3 = \Phi_0/4$, the interaction Hamiltonian vanishes, thus isolating the qubit from the bosonic field. State of the art technology [14] allows the variation of this external flux in times of about 0.1 ns, which is much faster than the interaction dynamics.

The proposed experiment of past-future entanglement extraction would constitute a fundamental proof for the possibility of using the vacuum state of a field to produce correlations between a qubit in the past and a qubit in the future, but this is not the only interesting aspect. From the quantum technologies point of view, the ability to extract past-future entanglement from the field using a pair of qubits could be used to implement a device which teleports a quantum state in time - as first suggested in [11] -, or in other words, we could use the field in the transmission line for building up a novel kind of quantum memory. In the following, we will explain this in more detail.

For the quantum teleportation in time, an observer - say, Paula - in possession of the qubit P and another qubit P' that she wants to teleport, should be able to carry out measurements on her qubits once the interaction had been disconnected at $-t_1$. At some time in the future after t_2, an observer - say, Frank - would use the results of Paula's measurements stored as classical information and locally manipulate the qubit F , in order to transfer the state of P' to F. The fidelity will be a function of the amount of quantum correlations between P and F. Note that during T_{off} the information of the quantum state of P' is in some sense codified in the field along the transmission line, even if we destroy P after its interaction and measurement. Then, this information is recovered and embodied in F after its interaction and the use of the stored classical bits.

The experimental realization of quantum teleportation has already been achieved in cQED with fairly high fidelity [24], up to the final step of single-shot qubit measurements. The fact that P and F are non-pure states should not prevent us from using them as a resource for teleportation, given the amount of quantum correlations both share [25, 26]. As shown in figure 2, entanglement is strong enough to consider the possibility of relatively high-fidelity teleportation for quantum states that are separated by times of nanoseconds. Since this interval is similar to the coherence times of the qubit, the scheme might be used as a quantum memory - provided that the coherence of the field is long enough -, therefore paving the way for future implementations where this time separation could be larger. In our setting, that time-lapse grows with the qubit spatial separation and the inverse of the qubit gap, those values being limiting experimental factors for it to work as a memory.

In a nutshell, we have proposed a experimentally feasible circuit QED setup to test the extraction of quantum correlations between different times contained in the vacuum of a quantum field. We have shown in particular that sizable past-future vacuum correlations can be transferred to a pair of qubits P and F, which only interact with the field in the past or the future respectively, even if the qubits do not coexist at the same time. Moreover, we discuss the possible technological uses of that entanglement extraction and the limitations of our scheme to work as a quantum memory.

The authors would like to acknowledge J. J. García-Ripoll, T. C. Ralph, A. Dragan, I. Fuentes, G. Adesso, J. Louko, N. Friis, C. M. Wilson, P. Forn-Díaz, and J. León for useful comments. This work is supported by Spanish MICINN Projects FIS2011-29287 and FIS2009-10061and CAM research consortium QUITEMAD S2009-ESP-1594. B. Peropadre and M. del Rey were supported by a CSIC JAE-PREDOC grant. M.del Rey was also supported by Residencia de Estudiantes.

---

[*] Electronic address: csl@iff.csic.es

[1] W. G. Unruh, Phys. Rev. D **14**, 870 (1976).

[2] S. J. Summers and R. F. Werner, Phys. Lett. A **110**, 257 (1985).

[3] S. J. Summers and R. F. Werner, J. Math. Phys. **28**, 2448 (1987).

[4] B. Reznik, A. Retzker, and J. Silman, Phys. Rev. A **71**, 042104 (2005).

[5] A. Retzker, J. I. Cirac, and B. Reznik, Phys. Rev. Lett. **94**, 050504 (2005).

[6] S. Marcovitch, A. Retzker, M. B. Plenio, and B. Reznik, Phys. Rev. A **80**, 012325 (2009).

[7] C. Sabín, J. J. García-Ripoll, E. Solano, and J. León, Phys. Rev. B **81**, 184501 (2010).

[8] M. Cliche and A. Kempf, Phys. Rev. A **81**, 012330 (2010).

[9] S. J. Olson and T. C. Ralph, Phys. Rev. Lett. **106**, 110404 (2011).

[10] Although in [9] these correlations are referred to as entanglement, such a wording might not be appropriate since quantum states in past and future belong to Hilbert spaces defined in different simultaneity planes. Nevertheless, quantum correlations of this sort can in principle be extracted to physical systems and exploited as a resource for quantum information tasks as in the case of typical entanglement.

[11] S. J. Olson and T. C. Ralph, Phys. Rev. A **85**, 012306 (2012).

[12] A. Dragan and I. Fuentes, ArXiv e-prints (2011), 1105.1192.

[13] J. Q. You and F. Nori, Nature **474**, 589 (2011).

[14] C. M. Wilson, G. Johansson, A. Pourkabirian, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, Nature **479**, 376 (2011).

[15] J. Bourassa, J. M. Gambetta, A. A. Abdumalikov, O. Astafiev, and Y. Nakamura, Phys. Rev. A **80**, 032109 (2009).

[16] P. Forn-Díaz, J. Lisenfeld, D. Marcos, J. J. García-Ripoll, E. Solano, C. J. P. M. Harmans, and J. E. Mooij, Physical Review Letters **105**, 237001 (2010), 1005.1559.

[17] T. Niemczyk, F. Deppe, H. Huebl, E. P. Menzel, F. Hocke, M. J. Schwarz, J. J. Garcia-Ripoll, D. Zueco, T. Hümmer, E. Solano, et al., Nature Physics **6**, 772 (2010).

[18] Note that we will use the term "past-future" correlations to refer to a different and more general notion than "timelike entanglement" or "entanglement between the past and the future light cones".

[19] DeWitt, General Relativity; an Einstein Centenary Survey (Cambridge University Press (Cambridge, UK), 1980), ISBN 0521299284.

[20] D. E. Bruschi, J. Louko, E. Martín-Martínez, A. Dragan, and I. Fuentes, Phys. Rev. A **82**, 042332 (2010).

[21] To have a mirror interpretation for a plane-wave-like basis like that provided in [9] we would need observers undergoing forbidden trajectories in the spacetime - horizontal hyperbolas in past and future light cones.

[22] B. Peropadre, P. Forn-Díaz, E. Solano, and J. J. García-Ripoll, Phys. Rev. Lett. **105**, 023601 (2010).

[23] M. Steffen, M. Ansmann, R. C. Bialczak, N. Katz, E. Lucero, R. McDermott, M. Neeley, E. M. Weig, A. N. Cleland, and J. M. Martinis, Science **313**, 1423 (2006).

[24] M. Baur, A. Fedorov, L. Steffen, S. Filipp, M. P. da Silva, and A. Wallraff, Phys. Rev. Lett. **108**, 040502 (2012).

[25] G. Bowen and S. Bose, Phys. Rev. Lett. **87**, 267901 (2001).

[26] S. Albeverio, S. M. Fei, and W. Yang, Commun. Theor. Phys. **38**, 301 (2002).